

SCHEMATIC DIAGRAM

Topic	Concepts	Degree of importance	References
			NCERT Text Book XII Ed. 2007
Relations & Functions	(i).Domain , Co domain & Range of a relation	*	(Previous Knowledge)
	(ii).Types of relations	***	Ex 1.1 Q.No- 5,9,12
	(iii).One-one , onto & inverse of a function	***	Ex 1.2 Q.No- 7,9
	(iv).Composition of function	*	Ex 1.3 QNo- 7,9,13
	(v).Binary Operations	***	Example 45 Ex 1.4 QNo- 5,11

SOME IMPORTANT RESULTS/CONCEPTS

** A **relation R** in a set A is called

(i) *reflexive*, if $(a, a) \in R$, for every $a \in A$,

(ii) *symmetric*, if $(a_1, a_2) \in R$ implies that $(a_2, a_1) \in R$, for all $a_1, a_2 \in A$.

(iii) *transitive*, if $(a_1, a_2) \in R$ and $(a_2, a_3) \in R$ implies that $(a_1, a_3) \in R$, for all $a_1, a_2, a_3 \in A$.

** Equivalence Relation : R is equivalence if it is reflexive, symmetric and transitive.

** **Function** :A relation $f : A \rightarrow B$ is said to be a function if every element of A is correlated to unique element in B.

* A is domain

* B is codomain

* For any x element $x \in A$, function f correlates it to an element in B, which is denoted by $f(x)$ and is called image of x under f . Again if $y = f(x)$, then x is called as pre-image of y .

* Range = $\{f(x) \mid x \in A\}$. Range \subseteq Codomain


* The largest possible domain of a function is called domain of definition.

** **Composite function** :

Let two functions be defined as $f : A \rightarrow B$ and $g : B \rightarrow C$. Then we can define a function

$\phi : A \rightarrow C$ by setting $\phi(x) = g\{f(x)\}$ where $x \in A$, $f(x) \in B$, $g\{f(x)\} \in C$. This function

$\phi : A \rightarrow C$ is called the composite function of f and g in that order and we write. $\phi = g \circ f$.



**** Different type of functions :** Let $f : A \rightarrow B$ be a function.

* f is **one to one (injective) mapping**, if any two different elements in A is always correlated to different elements in B , i.e. $x_1 \neq x_2 \Rightarrow f(x_1) \neq f(x_2)$ or, $f(x_1) = f(x_2) \Rightarrow x_1 = x_2$

* f is **many one mapping**, if \exists at least two elements in A such that their images are same.

* f is **onto mapping** (surjective), if each element in B is having at least one preimage.

* f is **into mapping** if $\text{range} \subseteq \text{codomain}$.

* f is **bijective mapping** if it is both *one to one and onto*.

**** Binary operation :** A binary operation $*$ on a set A is a function $*$: $A \times A \rightarrow A$. We denote $*(a, b)$ by $a * b$.

* A binary operation ' $*$ ' on A is a rule that associates with every ordered pair (a, b) of $A \times A$ a unique element $a * b$.

* An operation ' $*$ ' on A is said to be commutative iff $a * b = b * a \forall a, b \in A$.

* An operation ' $*$ ' on A is said to be associative iff $(a * b) * c = a * (b * c) \forall a, b, c \in A$.

* Given a binary operation $*$: $A \times A \rightarrow A$, an element $e \in A$, if it exists, is called **identity** for the operation $*$, if $a * e = a = e * a, \forall a \in A$.

* Given a binary operation $*$: $A \times A \rightarrow A$ with the identity element e in A , an element $a \in A$ is said to be **invertible** with respect to the operation $*$, if there exists an element b in A such that $a * b = e = b * a$ and b is called the **inverse of a** and is denoted by a^{-1} .

ASSIGNMENTS

(i) Domain , Co domain & Range of a relation

LEVEL I

1. If $A = \{1,2,3,4,5\}$, write the relation $a R b$ such that $a + b = 8, a, b \in A$. Write the domain, range & co-domain.

2. Define a relation R on the set \mathbf{N} of natural numbers by

$$R = \{(x, y) : y = x + 7, x \text{ is a natural number less than } 4 ; x, y \in \mathbf{N}\}.$$

Write down the domain and the range.

2. Types of relations

LEVEL II

1. Let R be the relation in the set \mathbf{N} given by $R = \{(a, b) | a = b - 2, b > 6\}$
Whether the relation is reflexive or not ? justify your answer.

2. Show that the relation R in the set \mathbf{N} given by $R = \{(a, b) | a \text{ is divisible by } b, a, b \in \mathbf{N}\}$ is reflexive and transitive but not symmetric.

3. Let R be the relation in the set \mathbf{N} given by $R = \{(a, b) | a > b\}$ Show that the relation is neither reflexive nor symmetric but transitive.

4. Let R be the relation on \mathbf{R} defined as $(a, b) \in R$ iff $1 + ab > 0 \forall a, b \in \mathbf{R}$.

(a) Show that R is symmetric.

(b) Show that R is reflexive.

(c) Show that R is not transitive.

5. Check whether the relation R is reflexive, symmetric and transitive.

$$R = \{(x, y) | x - 3y = 0\} \text{ on } A = \{1, 2, 3, \dots, 13, 14\}.$$

LEVEL III

1. Show that the relation R on A, $A = \{ x \mid x \in \mathbb{Z}, 0 \leq x \leq 12 \}$,
 $R = \{(a, b) : |a - b| \text{ is multiple of } 3\}$ is an equivalence relation.
2. Let N be the set of all natural numbers & R be the relation on $N \times N$ defined by
 $\{(a, b) R (c, d) \text{ iff } a + d = b + c\}$. Show that R is an equivalence relation.
3. Show that the relation R in the set A of all polygons as:
 $R = \{(P_1, P_2), P_1 \& P_2 \text{ have the same number of sides}\}$ is an equivalence relation. What is the set of all elements in A related to the right triangle T with sides 3, 4 & 5 ?
4. Show that the relation R on A, $A = \{ x \mid x \in \mathbb{Z}, 0 \leq x \leq 12 \}$,
 $R = \{(a, b) : |a - b| \text{ is multiple of } 3\}$ is an equivalence relation.
5. Let N be the set of all natural numbers & R be the relation on $N \times N$ defined by
 $\{(a, b) R (c, d) \text{ iff } a + d = b + c\}$. Show that R is an equivalence relation. [CBSE 2010]
6. Let A = Set of all triangles in a plane and R is defined by $R = \{(T_1, T_2) : T_1, T_2 \in A \& T_1 \sim T_2\}$
Show that the R is equivalence relation. Consider the right angled Δ s, T_1 with size 3, 4, 5;
 T_2 with size 5, 12, 13; T_3 with side 6, 8, 10; Which of the pairs are related?

(iii) One-one, onto & inverse of a function

LEVEL I

1. If $f(x) = x^2 - x^{-2}$, then find $f(1/x)$.
2. Show that the function $f: \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = x^2$ is neither one-one nor onto.
3. Show that the function $f: \mathbb{N} \rightarrow \mathbb{N}$ given by $f(x) = 2x$ is one-one but not onto.
4. Show that the signum function $f: \mathbb{R} \rightarrow \mathbb{R}$ given by: $f(x) = \begin{cases} 1, & \text{if } x > 0 \\ 0, & \text{if } x = 0 \\ -1, & \text{if } x < 0 \end{cases}$
is neither one-one nor onto.
5. Let $A = \{-1, 0, 1\}$ and $B = \{0, 1\}$. State whether the function $f: A \rightarrow B$ defined by $f(x) = x^2$ is bijective.
6. Let $f(x) = \frac{x-1}{x+1}$, $x \neq -1$, then find $f^{-1}(x)$

LEVEL II

1. Let $A = \{1, 2, 3\}$, $B = \{4, 5, 6, 7\}$ and let $f = \{(1, 4), (2, 5), (3, 6)\}$ be a function from A to B.
State whether f is one-one or not. [CBSE 2011]
2. If $f: \mathbb{R} \rightarrow \mathbb{R}$ defined as $f(x) = \frac{2x-7}{4}$ is an invertible function. Find $f^{-1}(x)$.
3. Write the number of all one-one functions on the set $A = \{a, b, c\}$ to itself.
4. Show that function $f: \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = 7 - 2x^3$ for all $x \in \mathbb{R}$ is bijective.
5. If $f: \mathbb{R} \rightarrow \mathbb{R}$ is defined by $f(x) = \frac{3x+5}{2}$. Find f^{-1} .

LEVEL III

1. Show that the function $f: \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = \frac{2x-1}{3}$, $x \in \mathbb{R}$ is one-one & onto function. Also find the f^{-1} .

2. Consider a function $f: \mathbb{R}_+ \rightarrow [-5, \infty)$ defined $f(x) = 9x^2 + 6x - 5$. Show that f is invertible & $f^{-1}(y) = \frac{\sqrt{y+6}-1}{3}$, where $\mathbb{R}_+ = (0, \infty)$.

3. Consider a function $f: \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = 4x + 3$. Show that f is invertible & $f^{-1}: \mathbb{R} \rightarrow \mathbb{R}$ with $f^{-1}(y) = \frac{y-3}{4}$.

4. Show that $f: \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = x^3 + 4$ is one-one, onto. Show that $f^{-1}(x) = (x-4)^{1/3}$.

5. Let $A = \mathbb{R} - \{3\}$ and $B = \mathbb{R} - \{1\}$. Consider the function $f: A \rightarrow B$ defined by

$$f(x) = \left(\frac{x-2}{x-3} \right). \text{ Show that } f \text{ is one one onto and hence find } f^{-1}. \quad [\text{CBSE2012}]$$

6. Show that $f: \mathbb{N} \rightarrow \mathbb{N}$ defined by $f(x) = \begin{cases} x+1, & \text{if } x \text{ is odd} \\ x-1, & \text{if } x \text{ is even} \end{cases}$ is both one one onto.

[CBSE2012]

(iv) Composition of functions

LEVEL I

1. If $f(x) = e^{2x}$ and $g(x) = \log \sqrt{x}$, $x > 0$, find

(a) $(f+g)(x)$ (b) $(f \cdot g)(x)$ (c) $f \circ g(x)$ (d) $g \circ f(x)$.

2. If $f(x) = \frac{x-1}{x+1}$, then show that (a) $f\left(\frac{1}{x}\right) = -f(x)$ (b) $f\left(-\frac{1}{x}\right) = \frac{-1}{f(x)}$

LEVEL II

1. Let $f, g: \mathbb{R} \rightarrow \mathbb{R}$ be defined by $f(x) = |x|$ & $g(x) = [x]$ where $[x]$ denotes the greatest integer function. Find $f \circ g(5/2)$ & $g \circ f(-\sqrt{2})$.

2. Let $f(x) = \frac{x-1}{x+1}$. Then find $f(f(x))$

3. If $y = f(x) = \frac{3x+4}{5x-3}$, then find $(f \circ f)(x)$ i.e. $f(y)$

4. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be defined as $f(x) = 10x + 7$. Find the function $g: \mathbb{R} \rightarrow \mathbb{R}$ such that $g \circ f(x) = f \circ g(x) = I_{\mathbb{R}}$ [CBSE2011]

5. If $f: \mathbb{R} \rightarrow \mathbb{R}$ be defined as $f(x) = (3-x^3)^{1/3}$, then find $f \circ f(x)$.

[CBSE2010]

6. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ & $g: \mathbb{R} \rightarrow \mathbb{R}$ be defined as $f(x) = x^2$, $g(x) = 2x - 3$. Find $f \circ g(x)$.

(v) Binary Operations

LEVEL I

1. Let $*$ be the binary operation on N given by $a*b = \text{LCM of } a \text{ \& } b$. Find $3*5$.
2. Let $*$ be the binary operation on N given by $a*b = \text{HCF of } \{a, b\}$, $a, b \in N$. Find $20*16$.
3. Let $*$ be a binary operation on the set Q of rational numbers defined as $a * b = \frac{ab}{5}$.

Write the identity of $*$, if any.

4. If a binary operation ' $*$ ' on the set of integer Z , is defined by $a * b = a + 3b^2$
Then find the value of $2 * 4$.

LEVEL 2

1. Let $A = N \times N$ & $*$ be the binary operation on A defined by $(a, b) * (c, d) = (a+c, b+d)$
Show that $*$ is (a) Commutative (b) Associative (c) Find identity for $*$ on A , if any.
2. Let $A = Q \times Q$. Let $*$ be a binary operation on A defined by $(a, b) * (c, d) = (ac, ad+b)$.
Find: (i) the identity element of A (ii) the invertible element of A .
3. Examine which of the following is a binary operation

(i) $a * b = \frac{a+b}{2}$; $a, b \in N$ (ii) $a * b = \frac{a+b}{2}$, $a, b \in Q$

For binary operation check commutative & associative law.

LEVEL 3

1. Let $A = N \times N$ & $*$ be a binary operation on A defined by $(a, b) * (c, d) = (ac, bd)$
 $\forall (a, b), (c, d) \in N \times N$ (i) Find $(2, 3) * (4, 1)$
(ii) Find $[(2, 3) * (4, 1)] * (3, 5)$ and $(2, 3) * [(4, 1) * (3, 5)]$ & show they are equal
(iii) Show that $*$ is commutative & associative on A .

2. Define a binary operation $*$ on the set $\{0, 1, 2, 3, 4, 5\}$ as $a * b = \begin{cases} a + b, & \text{if } a + b < 6 \\ a + b - 6, & \text{if } a + b \geq 6 \end{cases}$

Show that zero is the identity for this operation & each element of the set is invertible with $6 - a$ being the inverse of a .

[CBSE2011]

3. Consider the binary operations $*$: $\mathbf{R} \times \mathbf{R} \rightarrow \mathbf{R}$ and \circ : $\mathbf{R} \times \mathbf{R} \rightarrow \mathbf{R}$ defined as $a * b = |a - b|$ and $a \circ b = a$, $\forall a, b \in \mathbf{R}$. Show that $*$ is commutative but not associative, \circ is associative but not commutative.

[CBSE2012]

Questions for self evaluation

1. Show that the relation R in the set $A = \{1, 2, 3, 4, 5\}$ given by $R = \{(a, b) : |a - b| \text{ is even}\}$, is an equivalence relation. Show that all the elements of $\{1, 3, 5\}$ are related to each other and all the elements of $\{2, 4\}$ are related to each other. But no element of $\{1, 3, 5\}$ is related to any element of $\{2, 4\}$.
2. Show that each of the relation R in the set $A = \{x \in \mathbf{Z} : 0 \leq x \leq 12\}$, given by $R = \{(a, b) : |a - b| \text{ is a multiple of } 4\}$ is an equivalence relation. Find the set of all elements related to 1.

3. Show that the relation R defined in the set A of all triangles as $R = \{(T_1, T_2) : T_1 \text{ is similar to } T_2\}$, is equivalence relation. Consider three right angle triangles T_1 with sides 3, 4, 5, T_2 with sides 5, 12, 13 and T_3 with sides 6, 8, 10. Which triangles among T_1 , T_2 and T_3 are related?
4. If R_1 and R_2 are equivalence relations in a set A , show that $R_1 \cap R_2$ is also an equivalence relation.
5. Let $A = \mathbf{R} - \{3\}$ and $B = \mathbf{R} - \{1\}$. Consider the function $f : A \rightarrow B$ defined by $f(x) = \left(\frac{x-2}{x-3}\right)$.

Is f one-one and onto? Justify your answer.

6. Consider $f : \mathbf{R}^+ \rightarrow [-5, \infty)$ given by $f(x) = 9x^2 + 6x - 5$. Show that f is invertible and find f^{-1} .
7. On $\mathbf{R} - \{1\}$ a binary operation $'*'$ is defined as $a * b = a + b - ab$. Prove that $'*'$ is commutative and associative. Find the identity element for $'*'$. Also prove that every element of $\mathbf{R} - \{1\}$ is invertible.
8. If $A = \mathbf{Q} \times \mathbf{Q}$ and $'*'$ be a binary operation defined by $(a, b) * (c, d) = (ac, b + ad)$, for $(a, b), (c, d) \in A$. Then with respect to $'*'$ on A
- (i) examine whether $'*'$ is commutative & associative
 - (i) find the identity element in A ,
 - (ii) find the invertible elements of A .

ANSWERS

TOPIC 1 RELATIONS & FUNCTIONS

(i) Domain, Co domain & Range of a relation

LEVEL I

1. $R = \{ (3,5), (4,4), (5,3) \}$, Domain = $\{3, 4, 5\}$, Range = $\{3, 4, 5\}$

2. Domain = $\{1, 2, 3\}$, Range = $\{8, 9, 10\}$

(iii). One-one, onto & inverse of a function

LEVEL I

1. $-f(x)$ 6. $\frac{1+x}{1-x}$

LEVEL II

2. $f^{-1}(x) = \frac{(4x+7)}{2}$

3.6

5. $f^{-1}(x) = \frac{(2x-5)}{3}$

(iv). Composition of function

LEVEL II

5. $f \circ f(x) = x$

6. $4x^2 - 12x + 9$

(v) Binary Operations

LEVEL I

5. 15

2. 4

3. $e = 5$

4.50

Questions for self evaluation

2. $\{1, 5, 9\}$

3. T_1 is related to T_3

6. $f^{-1}(x) = \frac{\sqrt{x+6}-1}{3}$

7. $e = 0$, $a^{-1} = \frac{a}{a-1}$

8. Identity element $(1, 0)$, Inverse of (a, b) is $\left(\frac{1}{a}, \frac{-b}{a}\right)$