

chapter 1. Relations and Functions

Rvi Panchal

Relation

If A and B are two non-empty sets, then a relation R from A to B is a subset of $A \times B$.

If $R \subseteq A \times B$ and $(a, b) \in R$, then we say that a is related to b by the relation R , written as aRb .

Domain and Range of a Relation

Let R be a relation from a set A to set B . Then, set of all first components or coordinates of the ordered pairs belonging to R is called : the domain of R , while the set of all second components or coordinates = of the ordered pairs belonging to R is called the range of R .

Thus, domain of $R = \{a : (a, b) \in R\}$ and range of $R = \{b : (a, b) \in R\}$

Types of Relations

(i) **Void Relation** As $\Phi \subset A \times A$, for any set A , so Φ is a relation on A , called the empty or void relation.

(ii) **Universal Relation** Since, $A \times A \subseteq A \times A$, so $A \times A$ is a relation on A , called the universal relation.

(iii) **Identity Relation** The relation $I_A = \{(a, a) : a \in A\}$ is called the identity relation on A .

(iv) **Reflexive Relation** A relation R is said to be reflexive relation, if every element of A is related to itself.

Thus, $(a, a) \in R, \forall a \in A = R$ is reflexive.

(v) **Symmetric Relation** A relation R is said to be symmetric relation, iff

$(a, b) \in R (b, a) \in R, \forall a, b \in A$

i.e., $a R b \Rightarrow b R a, \forall a, b \in A$

$\Rightarrow R$ is symmetric.

(vi) **Anti-Symmetric Relation** A relation R is said to be anti-symmetric relation, iff

$(a, b) \in R$ and $(b, a) \in R \Rightarrow a = b, \forall a, b \in A$

(vii) Transitive Relation A relation R is said to be transitive relation, iff $(a, b) \in R$ and $(b, c) \in R$

$$\Rightarrow (a, c) \in R, \forall a, b, c \in A$$

(viii) Equivalence Relation A relation R is said to be an equivalence relation, if it is simultaneously reflexive, symmetric and transitive on A .

(ix) Partial Order Relation A relation R is said to be a partial order relation, if it is simultaneously reflexive, symmetric and anti-symmetric on A .

(x) Total Order Relation A relation R on a set A is said to be a total order relation on A , if R is a partial order relation on A .

Inverse Relation

If A and B are two non-empty sets and R be a relation from A to B , such that $R = \{(a, b) : a \in A, b \in B\}$, then the inverse of R , denoted by R^{-1} , is a relation from B to A and is defined by

$$R^{-1} = \{(b, a) : (a, b) \in R\}$$

Equivalence Classes of an Equivalence Relation

Let R be equivalence relation in A ($\neq \Phi$). Let $a \in A$.

Then, the equivalence class of a denoted by $[a]$ or $\{a\}$ is defined as the set of all those points of A which are related to a under the relation R .

Composition of Relation

Let R and S be two relations from sets A to B and B to C respectively, then we can define relation SoR from A to C such that $(a, c) \in SoR \Leftrightarrow \exists b \in B$ such that $(a, b) \in R$ and $(b, c) \in S$.

This relation SoR is called the composition of R and S .

(i) $RoS \neq SoR$

(ii) $(SoR)^{-1} = R^{-1}oS^{-1}$

known as **reversal rule**.

Rvi Panchal

Congruence Modulo m

Let m be an arbitrary but fixed integer. Two integers a and b are said to be congruence modulo m , if $a - b$ is divisible by m and we write $a \equiv b \pmod{m}$.

i.e., $a \equiv b \pmod{m} \Leftrightarrow a - b$ is divisible by m .

Important Results on Relation

- If R and S are two equivalence relations on a set A , then $R \cap S$ is also an equivalence relation on A .
- The union of two equivalence relations on a set is not necessarily an equivalence relation on the set.
- If R is an equivalence relation on a set A , then R^{-1} is also an equivalence relation on A .
- If a set A has n elements, then the number of reflexive relations from A to A is 2^{n^2-2} .
- Let A and B be two non-empty finite sets consisting of m and n elements, respectively. Then, $A \times B$ consists of mn ordered pairs. So, the total number of relations from A to B is 2^{nm} .

Binary Operations

Rvi Panchal

Closure Property

An operation $*$ on a non-empty set S is said to satisfy the closure property, if

$$a \in S, b \in S \Rightarrow a * b \in S, \forall a, b \in S$$

Also, in this case we say that S is closed for $*$.

An operation $*$ on a non-empty set S , satisfying the closure property is known as a binary operation.

or

Let S be a non-empty set. A function f from $S \times S$ to S is called a binary operation on S i.e., $f: S \times S \rightarrow S$ is a binary operation on set S .

Properties

- Generally binary operations are represented by the symbols $*$, $+$, \dots etc., instead of letters figure etc.
- Addition is a binary operation on each one of the sets N , Z , Q , R and C of natural numbers, integers, rationals, real and complex numbers, respectively. While addition on the set S of all irrationals is not a binary operation.
- Multiplication is a binary operation on each one of the sets N , Z , Q , R and C of natural numbers, integers, rationals, real and complex numbers, respectively. While multiplication on the set S of all irrationals is not a binary operation.
- Subtraction is a binary operation on each one of the sets Z , Q , R and C of integers, rationals, real and complex numbers, respectively. While subtraction on the set of natural numbers is not a binary operation.
- Let S be a non-empty set and $P(S)$ be its power set. Then, the union and intersection on $P(S)$ is a binary operation.

- Division is not a binary operation on any of the sets N, Z, Q, R and C . However, it is not a binary operation on the sets of all non-zero rational (real or complex) numbers.
- Exponential operation $(a, b) \rightarrow a^b$ is a binary operation on set N of natural numbers while it is not a binary operation on set Z of integers.

Types of Binary Operations

(i) Associative Law A binary operation $*$ on a non-empty set S is said to be associative, if $(a * b) * c = a * (b * c), \forall a, b, c \in S$.

Let R be the set of real numbers, then addition and multiplication on R satisfies the associative law.

(ii) Commutative Law A binary operation $*$ on a non-empty set S is said to be commutative, if

$$a * b = b * a, \forall a, b \in S.$$

Addition and multiplication are commutative binary operations on Z but subtraction not a commutative binary operation, since

$$2 - 3 \neq 3 - 2.$$

Union and intersection are commutative binary operations on the power $P(S)$ of all subsets of set S . But difference of sets is not a commutative binary operation on $P(S)$.

(iii) Distributive Law Let $*$ and \circ be two binary operations on a non-empty sets. We say that $*$ is distributed over \circ , if

$$a * (b \circ c) = (a * b) \circ (a * c), \forall a, b, c \in S \text{ also called (left distribution) and } (b \circ c) * a = (b * a) \circ (c * a), \forall a, b, c \in S \text{ also called (right distribution).}$$

Let R be the set of all real numbers, then multiplication distributes addition on R .

$$\text{Since, } a \cdot (b + c) = a \cdot b + a \cdot c, \forall a, b, c \in R.$$

(iv) Identity Element Let $*$ be a binary operation on a non-empty set S . An element $e \in S$, if it exist such that

$$a * e = e * a = a, \forall a \in S.$$

is called an identity elements of S , with respect to $*$.

For addition on R , zero is the identity elements in R .

$$\text{Since, } a + 0 = 0 + a = a, \forall a \in R$$

Rvi Panchal

For multiplication on \mathbb{R} , 1 is the identity element in \mathbb{R} .

Since, $a \times 1 = 1 \times a = a, \forall a \in \mathbb{R}$

Let $P(S)$ be the power set of a non-empty set S . Then, Φ is the identity element for union on $P(S)$ as

$A \cup \Phi = \Phi \cup A = A, \forall A \in P(S)$

Also, S is the identity element for intersection on $P(S)$.

Since, $A \cap S = S \cap A = A, \forall A \in P(S)$.

For addition on \mathbb{N} the identity element does not exist. But for multiplication on \mathbb{N} the identity element is 1.

(v) Inverse of an Element Let $*$ be a binary operation on a non-empty set ' S ' and let ' e ' be the identity element.

Let $a \in S$. we say that a^{-1} is invertible, if there exists an element $b \in S$ such that $a * b = b * a = e$

Also, in this case, b is called the inverse of a and we write, $a^{-1} = b$

Addition on \mathbb{N} has no identity element and accordingly \mathbb{N} has no invertible element.

Multiplication on \mathbb{N} has 1 as the identity element and no element other than 1 is invertible.

Let S be a finite set containing n elements. Then, the total number of binary operations on S is n^{n^2}

Let S be a finite set containing n elements. Then, the total number of commutative binary operation on S is $n \cdot [n(n+1)/2]$.

Rvi Panchal